

GALLAND/KIRBY ARTHROSCOPIC ECRB RELEASE / ELBOW ARTHROSCOPY POST-SURGICAL REHABILITATION PROTOCOL

POST-OP DAYS 1 – 7

- Dressing Posterior splint for 3-7 days; sutures out first week
- Shoulder, wrist, and hand active motion (AROM) Do not push into painful ROM
- Shoulder shrugs
- Ice as needed

GOAL

Pain control

WEEKS 1 - 2

- Splint D/C
- Counterforce Brace (elbow strap) worn with all exercises except stretches

- Continue appropriate previous exercises
- Elbow AROM
- Gentle Elbow / Wrist stretches
- · Gentle swimming / aqua-jogging
- Stationary bike

GOAL

Full elbow AROM

<u>WEEKS 2 - 6</u>

- Counterforce Brace Continue
- Continue appropriate previous exercises
- UBE Min resistance, gradually increase resistance and time as tolerated
- Isometrics x 8 (box plus supination / pronation) Pain-free, progress as tolerated
- Hand squeezing exercises Putty / sponge
- Treadmill Running progression program
- Elliptical trainer (light grip)

GOAL

Pain-free ADLs

<u>WEEKS 6 – 10</u>

- Counterforce Brace Continue
- Continue appropriate previous exercises
- PREs Wrist curls, reverse wrist curls, supination / pronation against resistance
- Pushup progression Wall to table to chair

GOAL

Normal elbow strength

<u>WEEKS 10 – 12</u>

- Counterforce brace Continue until 12 weeks post-op
- Continue appropriate previous exercises
- Pushups, regular
- · Weight training

GOAL • Return to full activities including sports

DEVELOPED BY:

Mark Galland, MD Kenneth Kirby, PT, DPT

ADAPTED FROM:

Physical Therapy Section William Beaumont Army Medical Center